

Bibliographie

1. Taheri S, Lin L, Austin D et al. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Med* 2004 ; 1 : e62.
2. Lacey JH, Stanley P, Hartmann M et al. The immediate effects of intravenous specific nutrients on EEG sleep. *Electroencephalogr Clin Neurophysiol* 1978 ; 44 : 275-80.
3. Afaghi A, O'Connor H, Chow CM. Acute effects of the very low carbohydrate diet on sleep indices. *Nutr Neurosci* 2008 ; 11 : 146-54.
4. Lacey JH, Crisp AH, Kalucy RS et al. Weight gain and the sleeping electroencephalogram: study of 10 patients with anorexia nervosa. *Br Med J* 1975 ; 4 : 556-8.
5. Karklin A, Driver HS, Buffenstein R. Restricted energy intake affects nocturnal body temperature and sleep patterns. *Am J Clin Nutr* 1994 ; 59 : 346-9.
6. Brondel L, Romer MA, Nougues PM et al. Acute partial sleep deprivation increases food intake in healthy men. *Am J Clin Nutr* 2010 ; 91 : 1550-9.
7. Bosy-Westphal A, Hinrichs S, Jauch-Chara K et al. Influence of partial sleep deprivation on energy balance and insulin sensitivity in healthy women. *Obes Facts* 2008 ; 1 : 266-73.
8. Tasali E, Leproult R, Spiegel K. Reduced sleep duration or quality: relationships with insulin resistance and type 2 diabetes. *Prog Cardiovasc Dis* 2009 ; 51 : 381-91.
9. Spiegel K, Tasali E, Penev P, Van Cauter E. Brief communication: Sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and appetite. *Ann Intern Med* 2004 ; 141 : 846-50.
10. Nedeltcheva AV, Kilkus JM, Imperial J et al. Sleep curtailment is accompanied by increased intake of calories from snacks. *Am J Clin Nutr* 2009 ; 89 : 126-33.
11. Nedeltcheva AV, Kilkus JM, Imperial J et al. Insufficient sleep undermines dietary efforts to reduce adiposity. *Ann Intern Med* 2010 ; 153 : 435-41.
12. Hartmann E. Effects of L-tryptophan on sleepiness and on sleep. *J Psychiatr Res* 1982-1983 ; 17 : 107-13.
13. Markus CR, Jonkman LM, Lammers JH et al. Evening intake of alpha-lactalbumin increases plasma tryptophan availability and improves morning alertness and brain measures of attention. *Am J Clin Nutr* 2005 ; 81 : 1026-33.
14. Wurtman RJ, Wurtman JJ, Regan MM et al. Effects of normal meals rich in carbohydrates or proteins on plasma tryptophan and tyrosine ratios. *Am J Clin Nutr* 2003 ; 77 : 128-32.
15. Herrera CP, Smith K, Atkinson F et al. High-glycaemic index and -glycaemic load meals increase the availability of tryptophan in healthy volunteers. *Br J Nutr* 2011 ; 105 : 1601-6.
16. Afaghi A, O'Connor H, Chow CM. High-glycemic-index carbohydrate meals shorten sleep onset. *Am J Clin Nutr* 2007 ; 85 : 426-30.
17. Bigard X, Simler N, Koulmann N. Nutrition, vigilance et sommeil : relations biologiques et comportementales. *Méd Somm* 2007 ; 4 : 15-23.
18. Bent S, Padula A, Moore D et al. Valerian for sleep: a systematic review and meta-analysis. *Am J Med* 2006 ; 119 : 1005-12.
19. Allen RP, Earley CJ. The role of iron in restless legs syndrome. *Mov Disord* 2007 ; 22 : S440-8.
20. Bonnet MH, Arand DL. Caffeine use as a model of acute and chronic insomnia. *Sleep* 1992 ; 15 : 526-36.
21. Feige B, Gann H, Brueck R. Effects of alcohol on polysomnographically recorded sleep in healthy subjects. *Alcohol Clin Exp Res* 2006 ; 30 : 1527-37.